


## REPORT DOCUMENTATION PAGE

FORM APPROVED  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of the collection of information, including suggestions for reducing the burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302 and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE		3. REPORT TYPE AND DATES COVERED Final June 1993 - May 1996	
4. TITLE AND SUBTITLE OF REPORT Metabolic and Enzymological Studies of Sulfur-Dependent Marine Hyperthermophiles				5. FUNDING NUMBERS (G) N00014-90-J-1894	
6. AUTHOR(S) Dr. Michael W. W. Adams					
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Georgia Research Foundation., Inc. 621 Boyd Graduate Research Center Athens, Georgia 30602				8. PERFORMING ORGANIZATION REPORT NUMBER: 10-21-RR166-301 10-21-RR166-333 10-21-RR166-352	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Dr. Eric Eisenstadt Office of Naval Research (1141-MB) 800 North Quincy Street Arlington, Virginia 22217				10. SPONSORING/MONITORING AGENCY REPORT NUMBER: N00014-90-J-1894	
11. SUPPLEMENTARY NOTES: Final Progress Report submitted to ONR on 9/30/96.					
12a. DISTRIBUTION AVAILABILITY STATEMENT 				12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) <p>The results from this study have provided the first insights into how hyperthermophilic organisms obtain energy from the metabolism of C and N compounds, and how this can be achieved at high temperatures. These organisms have unusual pathways based on the rarely-used element tungsten (W), which is present in the novel enzymes AOR, FOR and GAPOR. W is seldom used in biological systems, the analogous element, molybdenum (Mo), is virtually ubiquitous. We propose that W is much more suited to catalyze low potential reactions (such as those catalyzed by GAPOR, AOR and FOR) at extreme temperatures, and that such reactions could not be catalyzed by Mo-containing enzymes. We have cloned and sequenced the first genes for any W-protein, and the first genes for hyperthermophilic oxidoreductases. In addition, a new pathway for peptide metabolism has been proposed, again involving new types of enzyme, such as VOR, IOR and ACS. We also provided the first definitive model for the evolution of mesophilic oxidoreductases from hyperthermophilic enzymes, and the first crystal structure for a hyperthermophilic enzyme (AOR) was obtained, providing the first insights in to protein stability at extreme temperatures.</p>					
14. SUBJECT TERMS Hyperthermophiles, life at 100°C, archaea, tungsten-enzymes, metabolic enzymes, sulfur reduction.				15. NUMBER OF PAGES: 7	
				16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT:	18. SECURITY CLASSIFICATION OF THIS PAGE	19. SECURITY CLASSIFICATION OF ABSTRACT	20. LIMITATION OF ABSTRACT		

1785

# The University of Georgia

Department of Biochemistry & Molecular Biology

Dr. Eric Eisenstadt  
Office of Naval Research (1141-MB)  
800 North Quincy Street  
Arlington, Virginia 22217  
Tel: 703 696-4596

10-21-RR166-301  
10-21-RR166-333  
10-21-RR166-352

September 30, 1996

Dear Eric,


Please find attached two copies of my Final Progress Report for grant number N00014-90-J-1894 for the period June, 1993 - May, 1996.

I would like to thank both you and your office for supporting my research over this period. I feel we have been very productive and have made tremendous progress in understanding the metabolism and biochemistry of hyperthermophilic organisms.

Please do not hesitate to contact me if further information is required.

With best regards,

Yours sincerely,



Michael W. W. Adams  
Professor of Biochemistry, Molecular Biology and Microbiology

(Tel: 706 542-2060; FAX: 706 542-0229)  
(E-mail: ADAMSM@BSCR.UGA.EDU)

~~cc: Catherine Graddock, Sponsored Programs, UGA~~

19970827 142

## FINAL PROGRESS REPORT

GRANT#: N00014-90-J-1894

R&T CODE: 3412-085

PRINCIPAL INVESTIGATOR:

Michael W. W. Adams

INSTITUTION:

University of Georgia

GRANT TITLE:

"Metabolic and Enzymological Studies of Sulfur-Dependent Marine Hyperthermophiles"

AWARD PERIOD:

1 June 1993 - 31 May 1996

OBJECTIVES:

Hyperthermophiles are a recently discovered group of marine organisms that grow above 90°C. Virtually all are archaea. The majority are strictly anaerobic heterotrophs that utilize peptides and reduce elemental sulfur ( $S^0$ ) to  $H_2S$ . Our aims are to understand the novel biochemistry that must be required to sustain life near 100°C.

ACCOMPLISHMENTS:

At the outset of this research we had already established that the growth of heterotrophic hyperthermophiles is dependent upon the rarely-used element, tungsten (W) and a novel tungstopterin-containing enzyme, aldehyde ferredoxin oxidoreductase (AOR), was proposed to play a key role in its primary metabolism. In the present award period, we cloned and sequenced the gene for AOR from *Pyrococcus furiosus* (Tmax 105°C, Pf) (1) and in collaboration with Dr. Doug Rees (Caltech), a crystal structure of AOR at 2.3Å resolution was obtained (2). This was the first structure of a hyperthermophilic, a W-, or a pterin-containing enzyme. Hence, it gave important insights into the nature and role of pterin cofactors as well as into potential mechanisms of protein stability (2). In addition, W was shown to affect the metabolic patterns of Pf (3). AOR was also been purified from *Thermococcus* ES-1 (Tmax 95°C, ES-1), an organism that is obligately dependent upon elemental sulfur ( $S^0$ ) for growth (4). Moreover, from the related organism, *Thermococcus litoralis* (Tmax 98°C, Tl), a second type of W-containing aldehyde-oxidizing enzyme (FOR) was also purified (5), and its gene was cloned and sequenced (1). This enzyme had about 60% sequence similarity to Pf AOR showing that the two enzymes are closely related. We subsequently showed that, in addition to AOR and FOR, a third type of W-enzyme is present in these hyperthermophilic organisms. This was termed glyceraldehyde-3-phosphate oxidoreductase (GAPOR) and it was purified from *P. furiosus* (6). GAPOR was proposed to play a central role in a novel glycolytic pathway in these organisms (6) in which it replaces GAP dehydrogenase. Other glycolytic enzymes appear to be of the usual type. For example, we purified the enolase from Pf (7) and showed that it resembles analogous mesophilic enzymes except in thermal stability. In contrast to GAPOR, the other two tungstoenzymes, AOR and FOR, are proposed to function in peptide metabolism, as discussed below. However, we also showed that hyperthermophiles such as Pf are unique in that their growth is obligately dependent upon W (8). This cannot be replaced by

molybdenum, an analogous element that is utilized by virtually all other life forms (see 9, 10).

In the fermentation of peptides by the hyperthermophiles, we showed that the first step in the utilization of aromatic amino acids is catalyzed by transaminases (ATs). Both Tl and Pf were to contain two distinct types and these were purified (11,12). However, we also found that these organisms contain a new type of enzyme involved in the catabolism of aromatic amino acids which we term indolepyruvate oxidoreductase (IOR) (13). IOR produces aryl acetyl CoAs and CO<sub>2</sub> from the transaminated forms of the aromatic amino acids. Another key enzyme in peptide utilization, glutamate dehydrogenase (GDH), has also been purified from Tl (14) and the deep sea isolate ES-4 (15). Our investigations into the pathways of amino acid oxidation by these proteolytic hyperthermophiles also showed that in addition to IOR, they contain three types of 2-keto acid oxidoreductases and these have also been purified. While IOR is specific for aromatic keto acids (13) derived from the aromatic amino acids, another new enzyme which we term VOR uses keto acids derived from branched chain amino acids (16). The third enzyme is termed KGOR, and this is specific for 2-ketoglutarate (17). The fourth is POR, which mainly utilizes pyruvate, and this had already been purified from Pf in earlier studies. In the ONR-supported work, the genes for VOR and POR have been cloned and sequenced from *P. furiosus* and the POR genes have been sequenced from the bacterium *Thermotoga maritima* (Tmax 90°C: 18). From these data we have constructed an evolutionary model to show how the 2-keto acid oxidoreductases in mesophilic organisms arose from the gene fusion of ancestral genes now represented in the hyperthermophiles. Moreover, a unifying mechanism has been proposed to account for the diverse nature of these enzymes (10).

The fermentation of amino acids by these organisms is complicated by our finding that ES-1 expresses of an unusual alcohol dehydrogenase (ADH) which has been purified, and this reduces aldehydes to alcohols to dispose of excess reductant (19). Tl also contains a similar alcohol dehydrogenase (ADH) which was also purified (14). Both enzymes preferentially reduce aldehydes to alcohols. The question arose, therefore, as to the source of the aldehydes. We solved this by showing that POR, IOR and VOR not only oxidize 2-keto acids, they non-oxidatively decarboxylate them to the corresponding acid (20). Moreover, we also propose that the tungstoenzymes AOR and FOR have a similar role, that is, they function to oxidize such aldehydes to the corresponding acid (4, 10). The question also arose as the fate of the CoA-derivatives generated by POR, VOR, IOR and KGOR. We showed (21) that these are utilized by an enzyme known as acetyl CoA synthetase (ACS), which converts the CoA derivatives, ADP and phosphate to the acid and ATP. In fact, Pf has two ACS isoenzymes, wherein ACS I utilizes the products of the POR and VOR reactions, whereas only ACS II utilizes the products of the IOR reaction. Neither enzyme uses succinyl CoA, which is produced by KGOR (21). Hence, we have proposed a completely new pathway by which amino acids are converted to carboxylic acids with energy conservation by ACS (10).

We have also shown that S<sup>0</sup> reduction by hyperthermophiles such as Pf is an energy yielding reaction in Pf, in spite of

previous assumptions (3), and two  $S^0$ -reducing enzymes have been characterized: sulfhydrogenase (22) and a NADP-dependent sulfide dehydrogenase (SuDH) (23). We also found that SuDH functions as a ferredoxin:NADP oxidoreductase. This led to the elucidation of the pathway of electron transfer with the reconstitution of a system that would evolve  $H_2$  from pyruvate at  $90^\circ C$ . This comprised pyruvate oxidoreductase, ferredoxin, SuDH, NADP and hydrogenase (24). Hence, this shows how all of the reduced ferredoxin generated during the fermentation of both peptides and carbohydrates can be oxidized to produce  $H_2$ .

Finally, we also initiated a project to map by 2D-electrophoresis the proteins of *P. furiosus* (25) which will hopefully be of use when results from the genomic sequencing project of *P. furiosus* become available. In addition to refs. 9, 10, other aspects of the research presented here are summarized in the reviews given in refs. 26 - 31.

#### SIGNIFICANCE:

Our results have provided the first insights into how hyperthermophilic organisms obtain energy from the metabolism of C and N compounds, and how this can be achieved at high temperatures. These organisms have unusual pathways based on the rarely-used element tungsten (W), which is present in the novel enzymes AOR, FOR and GAPOR. W is seldom used in biological systems, the analogous element, molybdenum (Mo), is virtually ubiquitous. We propose that W is much more suited to catalyze low potential reactions (such as those catalyzed by GAPOR, AOR and FOR) at extreme temperatures, and that such reactions could not be catalyzed by Mo-containing enzymes (12). We have cloned and sequenced the first genes for any W-protein, and the first genes for hyperthermophilic oxidoreductases. In addition, a new pathway for peptide metabolism has been proposed, again involving new types of enzyme, such as VOR, IOR and ACS. We also provided the first definitive model for the evolution of mesophilic oxidoreductases from hyperthermophilic enzymes, and the first crystal structure for a hyperthermophilic enzyme (AOR) was obtained, providing the first insights in to protein stability at extreme temperatures.

#### PUBLICATIONS AND REPORTS ACKNOWLEDGING ONR-N00014-90-J-1894:

1. Kletzin, A., Mukund, S., Kelley-Crouse, T. L., Chan, M. K., Rees, D. C. and Adams, M. W. W. (1995) "Molecular characterization of tungsten-containing enzymes from hyperthermophilic archaea: aldehyde ferredoxin oxidoreductase from *Pyrococcus furiosus* and formaldehyde ferredoxin oxidoreductase from *Thermococcus litoralis*" *J. Bacteriol.* **177**, 4817-4819
2. Chan, M. K., Mukund, S., Kletzin, A., Adams, M. W. W. and Rees, D. C. (1995) "Structure of the hyperthermophilic tungstoenzyme aldehyde ferredoxin oxidoreductase" *Science* **267**, 1463-1469
3. Schicho, R. N., Ma, K., Adams, M. W. W. and Kelly, R. M. (1993) "Bioenergetics of sulfur reduction in the hyperthermophilic archaeon, *Pyrococcus furiosus*" *J. Bacteriol.* **175**, 1823-1830

4. Heider, J., Ma, K. and M. W. W. Adams. (1995) "Purification, characterization and metabolic function of aldehyde ferredoxin oxidoreductase from the hyperthermophilic and proteolytic archaeon, *Thermococcus* strain ES-1" *J. Bacteriol.* **177**, 4757-4764
5. Mukund, S. and Adams, M. W. W. (1993) "Characterization of a novel tungsten-containing formaldehyde ferredoxin oxidoreductase from the extremely thermophilic archaeon, *Thermococcus litoralis*. A role for tungsten in peptide catabolism" *J. Biol. Chem.* **268**, 13592-13600
6. Mukund, S. and Adams, M. W. W. (1995) "Glyceraldehyde-3-phosphate ferredoxin oxidoreductase, a novel tungsten-containing enzyme with a potential glycolytic role in the hyperthermophilic archaeon, *Pyrococcus furiosus*" *J. Biol. Chem.* **270**, 8389-8392
7. Peak, M. J., Peak, J. G., Stevens, F. J., Blamey, J., Mai, X., Zhou, Z. H. and Adams, M. W. W. (1994) "The hyperthermophilic glycolytic enzyme enolase in the archaeon, *Pyrococcus furiosus*: comparison with mesophilic enolases" *Arch. Biochem. Biophys.* **313**, 280-286
8. Mukund, S. and Adams, M. W. W. (1996) "Tungsten in the three tungstoenzymes of the hyperthermophilic archaeon *Pyrococcus furiosus* is not replaced by cell growth in the presence of vanadium or molybdenum" *J. Bacteriol.* **178**, 163-167
9. Kletzin, A. and Adams, M. W. W. (1996) "Tungsten in biology" *FEMS Microbiol. Rev.* **18**, 5-64
10. Adams, M. W. W. and Kletzin, A. (1996) "Oxidoreductase-Type Enzymes and Redox Proteins Involved in the Fermentative Metabolisms of Hyperthermophilic Archaea" *Adv. Prot. Chem.* **48**, 101-180
11. Andreotti, G., Cubellis, M. V., Nitti, G., Sannia, G., Mai, X., Marino, G. and Adams, M. W. W. (1994) "Characterization of aromatic aminotransferases in the hyperthermophilic archaeon *Thermococcus litoralis*" *Eur. J. Biochem.* **220**, 543-549
12. Andreotti, G., Cubellis, M. V., Nitti, G., Sannia, G., Mai, X., Adams, M. W. W. and Marino, G. (1994) "An extremely thermostable aromatic aminotransferase from the hyperthermophilic archaeon *Pyrococcus furiosus*" *Biochim. Biophys. Acta* **1247**, 90-96
13. Mai, X. and Adams, M. W. W. (1994) "Indolepyruvate ferredoxin oxidoreductase from the hyperthermophilic archaeon, *Pyrococcus furiosus*: a new enzyme involved in peptide fermentation" *J. Biol. Chem.* **269**, 16726-16732
14. Ma, K., Robb, F. T. and Adams, M. W. W. (1994) "Purification and characterization of NADP-specific alcohol dehydrogenase and NADP-specific glutamate dehydrogenase from the hyperthermophilic archaeon *Thermococcus litoralis*" *Appl. Environ. Microbiol.* **60**, 562-568
15. DiRuggiero, J., Robb, F. T., Jagus, R., Klump, H. K., Borges, K. M., Mai, X., Kessel, M. and Adams, M. W. W. (1993) "Characterization, cloning, and in vitro expression of an extremely thermostable glutamate dehydrogenase from the hyperthermophilic archaeon ES4" *J. Biol. Chem.* **268**, 17767-17774
16. Heider, J., Mai, X. and Adams, M. W. W. (1996) "Characterization of 2-ketoisovalerate ferredoxin oxidoreductase, a new and reversible coenzyme A-dependent



- enzyme involved in peptide fermentation by hyperthermophilic archaea. *J. Bacteriol.* **178**, 780-787
17. Mai, X. and Adams, M. W. W. (1996) "Characterization of a fourth type of 2-keto acid oxidizing enzyme from hyperthermophilic archaea: 2-ketoglutarate ferredoxin oxidoreductase from *Thermococcus litoralis*" *J. Bacteriol.* (in press, Oct. 1996 issue)
  18. Kletzin, A. and Adams, M. W. W. (1996) Pyruvate and 2-ketoisovalerate oxidoreducases from *Pyrococcus furiosus* and pyruvate oxidoreductase from *Thermotoga maritima*: molecular and phylogenetic characterization. *J. Bacteriol.* **178**, 248-257
  19. Ma, K., Loessner, H., Heider, J., Johnson, M. K. and Adams, M. W. W. (1995) "Effects of elemental sulfur on the metabolism of the deep sea hyperthermophilic archaeon, *Thermococcus* strain ES-1: Purification and characterization of a novel, sulfur-regulated, non-heme iron alcohol dehydrogenase" *J. Bacteriol.* **177**, 4748-4756
  20. Ma, K., Hutchins, A., Sung, S.-H. S. and Adams, M. W. W. "Pyruvate ferredoxin oxidoreductase from the hyperthermophilic archaeon, *Pyrococcus furiosus*, functions as a coenzyme A-dependent pyruvate decarboxylase" *Proc. Natl. Acad. Sci.* (revised version submitted 8/96)
  21. Mai, X. and Adams, M. W. W. "Purification and characterization of two reversible acyl-CoA synthetases (ADP-forming) from the hyperthermophilic archaeon *Pyrococcus furiosus*" *J. Bacteriol.* (in press, Oct. 1996 issue)
  22. Ma, K., Schicho, R. N., Kelly, R. M. and Adams, M. W. W. (1993) "Hydrogenase of the hyperthermophile, *Pyrococcus furiosus*, is an elemental sulfur reductase or sulfhydrogenase: evidence for a sulfur-reducing hydrogenase ancestor" *Proc. Natl. Acad. Sci. USA* **90**, 5341-5344
  23. Ma, K. and Adams, M. W. W. (1994) "Sulfide dehydrogenase from the hyperthermophilic archaeon, *Pyrococcus furiosus*: a new multifunctional enzyme involved in the reduction of elemental sulfur" *J. Bacteriol.* **176**, 6509-6517
  24. Ma, K., Zhou, Z. H. and Adams, M. W. W. (1994) "Hydrogen production from pyruvate by enzymes purified from the hyperthermophilic archaeon, *Pyrococcus furiosus*: a key role for NADPH" *FEMS Microbiol. Lett.* **122**, 263-266
  25. Giometti, C. S., Tollaksen, S. L., Mukund, S., Zhou, Z. H., Ma, K., Mai, X. and Adams, M. W. W. (1995) "Two-dimensional gel electrophoresis mapping of proteins isolated from the hyperthermophile *Pyrococcus furiosus*" *J. Chromatog.* **698**, 341-349

#### Invited Refereed Reviews:

26. Kelly, R. M., Baross, J. A. and Adams, M. W. W. (1994) "Biotechnology of life in boiling water" *Chem. Britain* **30**, 555-558
27. Adams, M. W. W. and Kelly, R. M. (1994) "Thermostability and thermoactivity of enzymes from hyperthermophilic archaea" *Biorg. Med. Chem.* **2**, 659-667
28. Kelly, R. M. and Adams, M. W. W. (1994) "Metabolism in hyperthermophilic microorganisms" *Antonie van Leeuwenhoek* **66**, 247-270

29. Adams, M. W. W. (1994) "Biochemical diversity among sulfur-dependent hyperthermophilic microorganisms" *FEMS Microbiol. Rev.* **15**, 261-277
30. Rees, D. C. and Adams, M. W. W. (1995) "Hyperthermophiles: taking the heat and loving it" *Structure* **15**, 251-254
31. Adams, M. W. W., Perler, F. B. and Kelly, R. M. (1995) "Extremozymes: expanding the limits of biocatalysis" *Bio/Technology* **13**, 662-668



## OBJECTIVES

### Hyperthermophilic Microorganisms Grow Near and Above 100°C:

- how do they stabilize proteins and cofactors at 100°C ?
- do they utilize conventional or novel primary metabolic pathways ?
- do they use conventional mechanisms for energy conservation ?

## ACCOMPLISHMENTS

- discovered that the growth of both carbohydrate- and peptide-fermenting "hyperthermophiles" is dependent upon tungsten (W), an element seldom used in biology.
- obtained the crystallographic structure of a novel tungstopterin-containing enzyme (AOR) at 2.3 Å resolution, the first for a W-, a pterin- or a hyperthermophilic enzyme.
- purified a third W-enzyme (GAPOR) which is involved in carbohydrate metabolism. This is in addition to AOR and FOR, which are involved in peptide catabolism.
- cloned and sequenced the first genes for any W-protein (AOR and FOR), and the first genes for any hyperthermophilic oxidoreductases.
- obtained the first definitive model for the evolution of mesophilic oxidoreductases from hyperthermophilic enzymes
- purified and characterized two more novel enzymes involved in peptide catabolism: ketoisovalerate oxidoreductase (VOR) and an unusual alcohol dehydrogenase.
- purified and characterized sulfide dehydrogenase, a novel enzyme involved in the reduction of elemental sulfur (S°), and defined the pathway for H<sub>2</sub> production from pyruvate.

## SIGNIFICANCE

- tungsten (W) is a key element in the primary metabolism of hyperthermophiles and first identification of a life form (the heterotrophic hyperthermophiles) that is obligately dependent upon W for growth
- first rationalization of W (but not the ubiquitous element, Mo) is essential for life near and above 100°C.
- several novel enzymes (including AOR, FOR, GAPOR, VOR, IOR, FNOR, ACS) have been discovered, purified and characterized
- proposed a new pathway of peptide fermentation and a modified glycolytic route
- first information on the mechanism of sulfur reduction by hyperthermophiles and the first characterization of enzymes that reduce S° at extreme temperatures.
- first definitive model for the evolution of mesophilic enzymes from enzymes of hyperthermophiles.
- first crystal structure of a hyperthermophilic or a W-containing enzyme.

M. W. W. Adams, U. of Georgia, Sept. 1996